

The 2012 North Carolina Agricultural Chemicals Manual is published by the North Carolina Cooperative Extension Service, College of Agriculture and Life Sciences, N.C. State University, Raleigh, N.C. These recommendations apply only to North Carolina. They may not be appropriate for conditions in other states and may not comply with laws and regulations outside North Carolina. These recommendations are current as of November 2011. Individuals who use agricultural chemicals are responsible for ensuring that the intended use complies with current regulations and conforms to the product label. Be sure to obtain current information about usage regulations and examine a current product label before applying any chemical. For assistance, contact your county Cooperative Extension agent. The use of brand names and any mention or listing of commercial products or services in this document does not imply endorsement by the North Carolina Cooperative Extension Service nor discrimination against similar products or services not mentioned.

BOOKMARKS

To jump to a location in this chapter without scrolling through the pages, use the Adobe Acrobat Bookmarks feature. To use Bookmarks, click on the Bookmarks icon on the left side of the page, then click on a Bookmarks entry.

IV — FERTILIZER USE

Lime and Fertilizer Suggestions for Field, Pasture, and Hay Crops	40
Fertilizer Suggestions for Tree Fruit	43
Fertilizer Suggestions for Small Fruit.....	44
Lime and Fertilizer Suggestions for Lawns	45
Fertilizer Suggestions for Ornamental Plants in Landscapes	48
Fertilizer Suggestions for Nursery Crops	48
Lime and Fertilizer Suggestions for Vegetable Crops	49
Fertilizer Rules and Regulations.....	53
Solubility of Selected Fertilizer Materials.....	56
Mixing Herbicides with Nitrogen Solutions or Fluid Fertilizers	57
How to Test Compatibility of Herbicides with Fluid Fertilizers.....	58
Fertilizer Placement.....	59
Livestock Manure Production Rates and Nutrient Content	59
Use of Municipal and Industrial Biosolids.....	63

Lime and Fertilizer Suggestions for Field, Pasture, and Hay Crops

**C. R. CROZIER and D. L. OSMOND, Soil Science; E. J. DUNPHY, L. FISHER, R. W. HEINIGER, D. L. JORDAN,
J. M. LUGINBUHL, P. MUELLER, and R. WEISZ, Crop Science; D. H. HARDY, NCDA&CS**

For best results, apply lime and nutrients (except nitrogen) based on soil test information. If this information is unavailable, use these suggestions as general guidelines. Ranges in rates are given for P_2O_5 and K_2O . Sandy soils generally respond to higher rates of K_2O than clayey ones, but the opposite is true for P_2O_5 . Also, attend to lime needs.

A liming program is essential to manage acidity in North Carolina soils. Base lime rates and frequency of applications on a routine soil sampling and testing program. In the absence of soil tests, most fields should receive 1 ton of lime per acre every 3 to 4 years. Liming more frequently at lesser rates may be necessary on sandy, light-colored soils where fertility changes more rapidly than in clay or organic soils, especially in years with excessive rainfall. Although liming is imperfect, over liming can cause micronutrient deficiencies, particularly manganese deficiency in small grain and soybeans grown on sandy soils. Dolomitic lime contains magnesium and is generally preferred over calcitic lime, especially on sandy soils. Lack of lime is the dominant soil fertility problem in North Carolina, even though liming is known to be a highly economical practice on either a short- or long-term basis.

Nitrogen rates are not based on soil testing. Use the RYE (Realistic Yield Expectation) for the soil and the appropriate nitrogen factor (NF) to establish the nitrogen rate. A range of NFs is found in Table 4-1A. The appropriate NF for each crop is determined by the soil mapping unit. Nitrogen rates based on RYE and NF are provided for each county by soil series at: <http://www.spatiallab.ncsu.edu/nutman/yields>.

TABLE 4-1A. LIME AND FERTILIZER SUGGESTIONS — FIELD CROPS

		Plant Nutrient Suggestions When Soil Test is Unavailable*			
Area of State or Soil Type	Optimum pH	N factor	P_2O_5 lb/acre	K_2O lb/acre	Remarks
CORN Grain					
Mineral soils	6.0	1.0 to 1.25 lb/bu	10 to 20	80 to 100	Banded starter fertilizer with 20 to 30 lb per acre of both N and P_2O_5 is recommended under no-till management or on cool, wet soils, especially when planting early. Starter P not likely to benefit on soils with very high initial P levels (P-index >100). Apply 1/4 to 1/3 of the N at planting. Sidedress remaining N when plants are 15 to 24 in. high. Under irrigation increase N rate by 10% to 15%. Starter band P if using no-till management or when soils are cool and wet. On deep sandy soils, apply K just before planting or use a split application of K at planting and at sidedress. Mineral soils that are sandy or greater than 18 in. to clay should receive 20 lb S per acre. Test organic soils to determine copper needs.
Mineral-organic soils	5.5				
Organic soils	5.0				
CORN Silage					
Mineral soils	6.0	10 to 12 lb/ton	10 to 20	100 to 120	
Mineral-organic soils	5.5				
Organic soils	5.0				
COTTON					
Mineral soils	6.2	0.06 to 0.12 lb/lb lint	10 to 20	50 to 70	Apply 20 to 30 lb N per acre at or before planting. Apply remaining N 2 to 3 weeks after first square. Starter P not likely to benefit on soils with very high initial P levels (P-index >100). After peanuts or soybeans, reduce total N by 25 to 30 lb per acre. Apply 0.5 lb per acre B at planting or as foliar spray at first bloom. On deep sands (>18 in. to clay), use higher end of lb per RYE range and apply 1/2 the remaining N at early square and 1/2 just prior to bloom. Deep sands should also receive 20 to 25 lb S per acre.
Mineral-organic soils	5.5	0.05 to 0.09			
Organic soils	5.0	0.03 to 0.06 lb/lb lint			
PEANUT					
Coastal plain	6.0	0	10 to 20	0	To minimize Ca deficiency risk, reduce unnecessary application of other cations. Apply K based on soil test recommendations and incorporate. During the growing season, at early to mid-flowering (late June-early July), apply gypsum to all Virginia market types. Use soil testing to determine Ca need for small-seeded runner market types. For larger-seeded "jumbo" runners, use half the rate as for Virginia market types. See Peanut Information 2012 (http://www.peanuts.ncsu.edu/) for more information on gypsum products and rates. Apply Mg only as recommended based on soil test. Apply 0.5 lb B per acre (liquid or dry). Apply Bradyrhizobia inoculant to seed or in the seed furrow regardless of previous rotation history to ensure peanut is capable of fixing N. Inoculation is especially important on land where peanut has not been grown recently. Apply 500 lb/ac of ammonium sulfate (i.e. 100 lb of N/ac) as soon as possible if roots are not nodulating effectively and foliage is N deficient. Zinc toxicity may occur in fields with soil test zinc indices greater than 250 or at lower indices with pH less than 6.0. Apply Mn at 1.0 lb Mn per acre when deficiency symptoms appear, often associated with high pH. See Peanut Information 2012 for selection of appropriate B and Mn products if needed.
SMALL GRAIN Grain					
Mineral soils	6.0	Wheat, Rye: 1.7 to 2.4 lb/bu Barley, Triticale: 1.4 to 1.6 lb/bu Oats: 1.0 to 1.3 lb/bu	10 to 20	80 to 100	Apply 15 to 20 lb N at planting (sometimes can be skipped if after soybean or peanut). If tiller density at spring green up (Feekes GS-3) is low, split N topdress (Feb., March). If tiller density is high, apply all N just before jointing. Use a tissue test at GS-5 to find the optimum nitrogen rate for wheat. Test organic soils to determine copper needs. Mineral soils that are sandy or greater than 18 in. to clay should receive 20 lb S per acre. Sensitive to manganese deficiency when soil pH is greater than 6.2.
Mineral-organic soils	5.5		10 to 20	60 to 80	
Organic soils	5.0		10 to 20	60 to 80	
SORGHUM Grain					
Mineral soils	6.0	1.5 to 2.0 lb/bu	10 to 20	50 to 70	Apply 20% to 25% of the N before planting. Apply remainder as a topdressing. If used for silage, increase N and K_2O by 40 lb per acre. Sensitive to manganese deficiency when soil pH is greater than 6.2.
Mineral-organic soils	5.5				
Organic soils	5.0				

TABLE 4-1A. LIME AND FERTILIZER SUGGESTIONS — FIELD CROPS

		Plant Nutrient Suggestions When Soil Test is Unavailable*			
Area of State or Soil Type	Optimum pH	N factor	P ₂ O ₅ lb/acre	K ₂ O lb/acre	Remarks
SOYBEAN					
Mineral soils	6.0	0	10 to 20	50 to 70	Fertilizer may be applied on preceding crop. Inoculate when planting in new land. If soil pH is low and lime is not applied, use 0.5 oz of sodium molybdate per acre seed treatment to facilitate N fixation (not a recommended substitute for lime). Test organic soils to determine copper needs. Soybean is sensitive to manganese deficiency when soil pH is greater than 6.2.
Mineral-organic soils	5.5				
Organic soils	5.0				
TOBACCO, BURLEY Greenhouse or Outdoor Float System - Float system: 5 to 7 days after seeding					
	5.5 to 6.0 (water)	See Remarks	See Remarks	See Remarks	75 to 100 ppm N from 20-10-20 or similar ratio fertilizer. Choose fertilizers with no more than 0.2% boron. Avoid fertilizers with 50% or more of N from urea. Test source water before using, and solutions during the season for nutrient levels, alkalinity, and conductivity.
TOBACCO, BURLEY Greenhouse or Outdoor Float System - 4 weeks after seeding					
	5.5 to 6.0 (water)	See Remarks	See Remarks	See Remarks	75 to 100 ppm N from 20-10-20 or similar ratio fertilizer. Test solutions for nutrient concentrations.
TOBACCO, BURLEY Field - Planting					
	6.0 (5.5 if history of Black Shank)	See Remarks	10 to 20	40 to 200	Apply 80 to 100 lb N per acre under conditions where no manure or legume is involved. Avoid excess chloride by using only fertilizers formulated for tobacco.
TOBACCO, BURLEY Field - Sidedressing					
	6.0 (5.5 if history of Black Shank)	See Remarks	0	0	Apply 100 to 150 lb N per acre 2 to 3 weeks after transplanting. It may contain up to 75% ammonium-N. For most soils, total N application (planting + sidedressing) of 180 to 200 lb per acre is adequate for optimum yield in the mountains and 250 lb is adequate in the piedmont.
TOBACCO, FLUE-CURED Greenhouse - Float system: 5 to 7 days after seeding					
Coastal plain and piedmont	5.5 to 6.0 (water)	See Remarks	See Remarks	See Remarks	100 to 150 ppm N from 20-10-20 or similar ratio fertilizer. Choose fertilizer with no more than 0.2% boron. Avoid fertilizers with 50% or more of N from urea. Test source water before using, and solutions during the season for nutrient levels, alkalinity, and conductivity.
TOBACCO, FLUE-CURED Greenhouse - Float system: 4 weeks after seeding					
Coastal plain and piedmont	5.5 to 6.0 (water)	See Remarks	See Remarks	See Remarks	100 ppm N from 20-10-20 or similar ratio fertilizer or ammonium nitrate. Test solutions for nutrient concentrations.
TOBACCO, FLUE-CURED Field - Planting					
Coastal plain and piedmont	6.0	See Remarks and <i>Flue-Cured Tobacco Information, AG-187</i>	10 to 20	90 to 110	Apply 35 to 40 lb N per acre from tobacco fertilizer containing up to 75% ammonium-N. If needed, apply P ₂ O ₅ at or within 7 days after transplanting. Piedmont soils are more likely to require fertilizer P than those in the coastal plain.
TOBACCO, FLUE-CURED Field - Sidedressing					
0 to 10 in. to clay		See Remarks	0	See Remarks	Apply 20 lb N per acre 2 to 3 weeks after transplanting. N source may contain up to 75% ammonium-N. N-P-K ratios of 1-0-0 or 1-0-1 are sufficient for most tobacco soils if 90 to 110 lb of K ₂ O per acre were supplied by the base fertilizer at planting. Use plant tissue analysis to identify nutrient deficiencies that need to be corrected.
11 to 15 in. to clay		See Remarks	0	See Remarks	Apply 30 lb N per acre under conditions described in remarks for 0 to 10 in. to clay. Use plant tissue analysis to identify nutrient deficiencies that need to be corrected.
Over 15 in. to clay		See Remarks and <i>Flue-Cured Tobacco Information, AG-187</i>	0	See Remarks	Apply 40 lb N per acre under conditions described in remarks for 0 to 10 in. to clay. Use plant tissue analysis to identify nutrient deficiencies that need to be corrected.
TOBACCO, FLUE-CURED Field - Adjustment for leaching					
		See Remarks and <i>Flue-Cured Tobacco Information, AG-187</i>			Replace N and K that are lost by leaching as early as conditions permit. Adjustments are normally not needed on soils with clay less than 10 in. from the surface. Magnesium and sulfur leaching may also be a concern on deep sandy soils. Use plant tissue analysis as a guide for determining nutrient deficiencies that need to be corrected.

TABLE 4-1B. LIME AND FERTILIZER SUGGESTIONS — PASTURE AND HAY CROPS

Commodity	Purpose	Area of State	Optimum pH	Plant Nutrient Suggestions When Soil Test is Unavailable*			Remarks
				N Based on RYE†	P ₂ O ₅ lb/acre	K ₂ O lb/acre	
ALFALFA	Seeding	All	6.5	See Remarks	90	150	Apply 20 lb N per acre before planting. Apply 3 lb boron per acre. Use inoculant at seeding.
	Annual maintenance	All	6.5	0	45	130	Apply first topdressing in spring of second year, following seeding. Each annual topdressing should include 2 lb boron per acre.
LADINO CLOVER/GRASS MIXTURE (>30% clover)	Seeding	All	6.5	See Remarks	90	150	Apply 20 lb N per acre before planting.
	Annual maintenance	All	6.5	0	45	150	Apply first topdressing in spring of second year, following seeding. When legumes make up 30% of stands, yield will be similar to pure grass receiving 150 to 200 lb of N per acre.
BLUEGRASS/WHITE CLOVER MIXTURE (>30% clover)	Annual maintenance	Mountain	6.0	0	40	90	
TALL FESCUE ORCHARD-GRASS TIMOTHY, PRAIRIEGRASS	Seeding	All	6.5	See Remarks	45	70	Apply 40 to 60 lb N per acre at planting.
	Annual maintenance	All	6.0	40 to 50 lb/ton‡	40	90	Use higher N rate for maximum production. Apply 1/2 N in February to March and 1/2 N in August to September; or 1/3 N in February, 1/3 N in April, and 1/3 N in August.
BERMUDA-GRASS, hybrid and improved seed cultivars	Sprigging	All	6.0	See Remarks	45	90	Apply 60 to 80 lb N per acre before planting. If sprigged in March, apply 1/2 N in May and 1/2 N in July first year. May apply additional 60 to 80 lb N/acre if complete soil covered before Sept. 1. If planted with a companion legume, lime to soil pH 6.5.
	Annual maintenance	All	6.0	40 to 50 lb/ton‡	40	180	Apply N in three or four applications, but not later than Sept. 1. If planted with a companion legume, lime to soil pH 6.5.
SUDAN HYBRID SORGHUM — Sudan hybrids PEARL MILLET RYEGRASS SMALL GRAIN	Seeding	All	6.0	See Remarks	40	90	Apply 50 to 60 lb N per acre planting. Apply N in two to four applications following each harvest except the last one.
	Topdressing	All	6.0	40 to 60 lb/ton‡	0	0	Apply N in two to four applications following each harvest except the last one.
RED CLOVER, FESCUE, ORCHARD-GRASS MIXTURE (>30% clover)	Seeding	Piedmont	6.5	See Remarks	90	150	Apply 20 lb N per acre before planting.
	Annual maintenance	Piedmont	6.5	0	25 to 35	60 to 80	Apply in second and third year if stand is adequate.
SWITCHGRASS FLACCIDGRASS CAUCASIAN BLUESTEM EASTERN GAMMAGRASS	Seeding	All	5.5 to 6.0	See Remarks	20	80	Apply 40 lb N per acre after plants are 6 to 10 in. high to minimize grassy weed competition.
	Maintenance	All	5.5 to 6.0	Switchgrass: 30 to 40 lb/ton All others: 35 to 45 lb/ton	20	80	Apply 1/2 N in May, 1/2 N in July, or 1/3 each in May, June, and July.

* Suggested rates may overestimate or underestimate actual needs. Take soil samples to assure accurate nutrient requirements.
 ‡ Rate is for tons of dry matter produced. Recognize that grazing animals excrete more than 80% of the nutrients they consume from feed; therefore, consider how grazing animals are managed to uniformly distribute the excreta over the pasture. To minimize N loss to the environment, N requirements for pastures should be reduced by 25% to 50% compared to what would be used for hay-silage production; yields may or may not be reduced.

Fertilizer Suggestions for Tree Fruit

M. L. PARKER, Horticultural Science; R. J. GEHL, Soil Science, and D. H. HARDY, NCDA&CS

TABLE 4-2. FERTILIZER SUGGESTIONS FOR TREE FRUIT

Purpose	Material	Amount	Precautions and Remarks
APPLES			
Preplant	Lime and P fertilizer	Depends on soil test.	Prepare soil as deep as possible before planting. Take soil samples at least 12 in. deep (preferably 0 to 8 in. and 8 to 16 in.) for lime and phosphorus recommendation. Apply one-half of total, adjusted for the depth of incorporation, and plow down; apply other half and work in well.
Improve and/or maintain growth of young trees	10-10-10 or its equivalent	1 lb for each year of age until tree begins bearing. Then as recommended from leaf analysis.	Apply before rainfall or irrigate after application before buds swell in the spring.
To raise boron level of tree	Solubar	1 lb/100 gal of spray at first cover	If leaf analysis shows a deficiency, use additional cover sprays as recommended from leaf analysis to reduce "cork spot." Dry years and large fruit may enhance the incidence of "cork spot."
Growth and fruit development	Nitrogen	1.25 lb of actual N for trees producing 10 to 15 bu of apples	Annual terminal growth should be about 12 in. Use observations of growth, crop size, and fruit condition plus the leaf analysis to determine the yearly application.
Increase foliar level	Potassium	Apply according to leaf analysis. Rate dependent upon soil analysis.	Leaf analysis is a good indicator of the need for a soil application.
Increase calcium level of tree	Gypsum (CaSO ₄)	15 to 50 lb/tree with a 6- to 10-ft radius.	Apply only as needed by low soil or tissue calcium. One application will usually last 3 to 5 years.
	Calcium nitrate	Apply in late fall or early spring at rate to supply recommended nitrogen.	Applied as soil applications to increase calcium supply and reduce "Bitter Pit."
Foliar application	Calcium nitrate	3 lb/100 gal for sprays two weeks apart and ending 2 to 3 weeks before harvest.	Apply to reduce the incidence of "Bitter Pit." Excessive use of CaNO ₃ may result in excessive tree vigor, which may actually worsen Bitter Pit. Both soil and foliar applications may be needed on large-fruited varieties. Leaf analysis may be beneficial.
	Calcium chloride	Same time as calcium nitrate. 2 lb/100 gal water.	Apply to reduce incidence of "Bitter Pit." DO NOT apply when temperature is 85°F or above.
PEACHES			
Preplant	Lime & Phosphorus	Depends on soil test.	Apply dolomitic lime necessary to raise soil pH above 6.0. Apply phosphorus to raise levels to desired range as indicated by soil test.
Tree growth, first year	N, P ₂ O ₅ , & K ₂ O	5 lb/acre of each per application.	Broadcast 0.5 lb of 10-10-10 around trees after growth starts in spring (April). Repeat every 4 to 6 weeks until August on sandy soils. On heavier soils, apply 0.5 lb of 10-10-10 one month after planting and 0.5 lb of 10-10-10 in May.
Tree growth, second year	N, P ₂ O ₅ , and K ₂ O	10 lb/acre of each per application.	Double amounts used first year. Make first application before growth starts in March and repeat in May. In sandy soils or if leaching is severe, an additional application may be made in July.
Tree growth, third year	N, P ₂ O ₅ , and K ₂ O	30 lb/acre of each	Make first application of 15 lb of each before growth starts and repeat in 6 to 8 weeks. If leaching is severe, repeat in July.
Growth and fruit development of young, bearing trees	N, P ₂ O ₅ , & K ₂ O	70 lb/acre of each (Determine by soil and foliar analysis.)	Broadcast under trees 40 lb/acre each of N, P ₂ O ₅ , and K ₂ O (for example 400 lb/acre 10-10-10) before growth starts. Add 30 lb/acre each of N, P ₂ O ₅ , and K ₂ O after fruit set. If soil phosphorus test is high, omit P in second application. If leaching is severe, apply 20 to 30 lb N/acre after harvest.
Growth and fruit development of mature trees	N, P ₂ O ₅ , and K ₂ O	70 lb/acre of each (Determine by soil and foliar analysis.)	Broadcast under trees 40 lb/acre each of N, P ₂ O ₅ , and K ₂ O (for example, 400 lb/acre 10-10-10) before growth starts. Add 30 lb/acre of N, P ₂ O ₅ , and K ₂ O after fruit set. If soil test phosphorus is high, omit P in second application. If leaching is severe, apply 30 lb N/acre after harvest.
Increase boron level of tree	Solubar		Apply boron to producing trees by spraying foliage with Solubar or equivalent product at a rate of 0.75 lb Solubar/100 gal of water once each year.
Soil pH maintenance	Dolomitic lime		Maintain soil pH above 6.0.

Fertilizer Suggestions for Small Fruit

BARCLAY POLING, B. CLINE, and G. E. FERNANDEZ, Horticultural Science; and B. R. CLEVELAND, NCDA&CS

For best results, fertilize using soil and tissue test information. If unavailable, use the general suggestions below.

TABLE 4-3. FERTILIZER SUGGESTIONS FOR SMALL FRUIT

Purpose	Material	Amount	Precautions and Remarks
BLACKBERRIES			
Preplant	Lime, P ₂ O ₅ , and K ₂ O	Depends on soil test.	Blackberries can be grown on a variety of soil types. Regardless of the soil type, however, organic matter additions, pH adjustments, and incorporation of phosphorus (P) and potassium (K) should be completed before planting to optimize productivity. Take a soil test three to six months prior to planting to ensure that the right soil amendments are added at the most efficient rates.
Growth first year	N	20 to 50 lb/acre	Fertilize within 1-2 weeks after an early spring planting and again 30 days later. Optimally, N can be portioned out through the drip irrigation system on a weekly basis.
	P ₂ O ₅ and K ₂ O	30 to 60 lb/acre	Assuming preplant P and K were applied according to soil test recommendations, additional P and K during the first year should not be needed. If this is not the case, apply P and K in 4-inch bands around but not closer than 6 inches from stems. Optimally, portion out P and K through the drip irrigation system on a weekly basis during the growing season.
Growth second year	N	35 to 60 lb/acre (up to 65 lb/acre on sandy soil)	Apply N in single or split application. For single application, apply in March. Spread fertilizer uniformly in 4-inch bands around but not closer than 6 inches from stems. For split application, apply N between the spring flush of growth and harvest with the greater portion applied as spring growth begins. Optimally, N can be portioned out through the drip irrigation system on a weekly basis. Only apply N after harvest if tissue analysis indicates it is low. Use primocane leaf tissue analysis 10-14 days post harvest to optimize N fertilization. Monitor growth and adjust N appropriately to achieve optimal growth rate; check excessive vegetative growth with a reduction in N rate.
	P ₂ O ₅ and K ₂ O	30 to 60 lb/acre	P and K can be applied in the fall if this is most convenient. Optimally, apply P and K during the growing season as per N recommendations. Adjust P and K rates as needed according to tissue analysis.
Growth third year and mature planting	N	60 to 80 lb/acre	Apply N, P and K according to recommendations for second year. Use primocane leaf tissue analysis 10-14 days post harvest to optimize and manage fertilization program.
	P ₂ O ₅ and K ₂ O	30 to 60 lb/acre	
BLUEBERRIES			
Growth first year	N, P ₂ O ₅ , and K ₂ O	40 to 80 lb 14-28-14/acre per application	Apply after first flush of growth and repeat every 4 to 6 weeks until mid-August. Extend application interval during dry periods until rainfall has totaled 4 inches Based on 1,360 plants per acre.
Growth second year	N, P ₂ O ₅ , and K ₂ O	Double first year amount at first application only	Use same schedule and amounts after first application as first year.
Growth and fruit development of bearing plants	N, P ₂ O ₅ , and K ₂ O	150 to 250 lb 14-28-14/acre	Apply 2/3 of this amount before bloom and 1/3 4 to 6 weeks later (early May). Apply 10 to 30 lb additional N/acre immediately after harvest if more vigorous growth is desired. Apply 50 lb per acre of diammonium phosphate (18-48-0) in mid-August to maintain plant vigor if P ₂ O ₅ is low or leaching has been severe.
GRAPES, BUNCH			
			Same as muscadine schedule. Petiole analysis can show which nutrients are limiting. Collect petioles opposite the first or second flower/fruit cluster at full bloom to veraiso. Contact your county Cooperative Extension agent for further information.
GRAPES, MUSCADINE			
Preplant	Lime, P ₂ O ₅ , and K ₂ O	Depends on soil test.	Test before planting and apply P ₂ O ₅ , K ₂ O, and lime according to soil test.
First year	N, P ₂ O ₅ , and K ₂ O	0.25 lb 10-10-10 per vine per application	Apply after growth starts (late April to early May) and repeat in June and July (but no later than mid-July as winter injury may occur). Broadcast in a circle at least 18 inches from the trunk.
Second year	N, P ₂ O ₅ , and K ₂ O	0.5 lb 10-10-10 per vine per application	Apply in March and again in May and early July. To minimize the potential for winter cold injury, piedmont and foothills growers should omit the July fertilizer application. Do not put fertilizer closer than 21 inches from the trunk.
Third year	N, P ₂ O ₅ , and K ₂ O	0.75 lb 10-10-10 per vine per application	Apply in March, in May and again in late June. Piedmont and foothills growers should omit the late June fertilizer application. Do not put fertilizer closer than 21 inches from the trunk.
Mature vines	N, P ₂ O ₅ , and K ₂ O	200 lb 10-10-10/acre per application	Apply in March (near bud break), and again in late May. If more vigorous growth is desired, add an additional 20 lb N per acre in late June (from 200 lb 10-10-10). Omit this last application in the piedmont and foothills. In Eastern NC, an alternative fertilizer to 10-10-10 that shows promise involves the application of 6-6-18 in March and mid-to-late May at the rate of 333 lb/acre per application (instead of 10-10-10 at 200 lb/acre per application). A final application of calcium nitrate is applied in late June at 133 lb /acre (provides 20 lb N per acre). Take leaf samples in early to mid-June to determine the actual nutritional status of mature vines. Collect a double fist full of mature leaves located opposite fruit clusters on fruiting shoots. Detach the petioles from the leaves before placing the leaf blades in a paper bag. Send samples to the Agronomic Division, NC Dept. of Agriculture & Consumer Services (see Chapter 3).
	B	1 lb Solubor/acre	For mature vineyards, a common recommendation has been to apply 1 lb of Solubor (20% boron) annually with 100 gallons of water per acre just before bloom. Boron deficiency is more likely on sandy soils with high pH. Excessive boron causes injury; do not exceed boron recommendations.
RASPBERRIES			
Preplant	Lime, P ₂ O ₅ , and K ₂ O	Depends on soil test	Test before planting and apply P ₂ O ₅ , and K ₂ O and lime according to soil test.
Growth first year	N, P ₂ O ₅ , and K ₂ O	250 to 500 lb10-10-10/acre	Fertilize 30 to 60 days after planting. Apply fertilizer in a band at the side of the row but not closer than 6 inches from stems. If using a drip system, the nutrients can be added via the drip system. Portion out the fertilizer at the recommended rates weekly or as needed.
Growth second year	N, P ₂ O ₅ , and K ₂ O	350 to 500 lb10-10-10/acre	Apply fertilizer in a band at the side of the row but not closer than 6 in. from stems. If using a drip system, the nutrients can be added via the drip system. Portion out the fertilizer at the recommended rates weekly or as needed.
Growth third year and mature planting	N, P ₂ O ₅ , and K ₂ O	500 to 800 lb10-10-10/acre	Apply fertilizer in a band at the side of the row but not closer than 6 in. from stems. If using a drip system, the nutrients can be added via the drip system. Portion out the fertilizer at the recommended rates weekly or as needed.

TABLE 4-3. FERTILIZER SUGGESTIONS FOR SMALL FRUIT

Purpose	Material	Amount	Precautions and Remarks
STRAWBERRIES, Matted-row			
Growth of new planting	N	30 to 40 lb N/acre	Apply in May and repeat in August or September on sandy soils. An additional 20 to 30 lb N per acre may be applied in January.
	P ₂ O ₅ , K ₂ O, and lime	Depends on soil test	Test before planting and apply P ₂ O ₅ , K ₂ O, and lime based on soil test.
Growth and fruit development	N	30 to 40 lb N/acre	Apply in August or September and in sandy soils again in January.
	N, P ₂ O ₅ , and K ₂ O	300 to 400 lb 10-10-10	Apply after harvest. If soil test for P and K are high, 30 to 40 lb of N may be used rather than 10-10-10.
STRAWBERRIES, Plasticulture			
Preplant (fall)	N	60 lb/acre	Broadcast and incorporate before bedding. Ammonium nitrate or a complete fertilizer (if P and K recommended) may be used.
	P ₂ O ₅ , K ₂ O, and lime	Refer to soil test. If not available, apply 60 lb P ₂ O ₅ and 120 lb K ₂ O/acre.	Soil test before planting. Broadcast and incorporate before bedding. Apply lime 3 months before planting.
Preharvest (spring)	N	1/2 to 1 lb/acre/day (3.5 to 7 lb/acre/week); depending on petiole nitrate test	Begin biweekly tissue testing when plants begin growing in the spring. Adjust rate or omit applications depending on tissue test interpretation. Another option is to apply recommended N in three or four applications over a 12-week period (3- or 4-week interval) beginning with first growth.
	Other nutrients	Depends on tissue test	Biweekly tissue tests will indicate need. If B is needed, apply at 1/8 lb B per acre.

Lime and Fertilizer Suggestions for Lawns

CHARLES PEACOCK, GRADY MILLER, and MATT MARTIN, Crop Science

SUGGESTED ESTABLISHMENT FERTILIZATION

Collect a soil sample for NCDA&CS analysis and follow lime and fertilizer recommendations. If the soil has not been tested, incorporate 75 pounds of ground limestone, except for centipedegrass, and 15 pounds of 0-14-14 fertilizer (or equivalent) per 1,000 square feet into the soil to a depth of 4 to 8 inches before seeding. At seeding, apply 1 pound nitrogen per 1,000 square feet from a turf-grade fertilizer in which one-fourth to one-half of the nitrogen is slowly available (e.g., 12-4-8 or 16-4-8). Use half of this fertilization rate when establishing centipedegrass. For more information, see *Carolina Lawns* or the lawn maintenance calendar for your specific grass. These can be found at <http://www.turfiles.ncsu.edu>. You can also request copies at your county Extension center.

TABLE 4-4A. SUGGESTED MAINTENANCE FERTILIZATION FOR COASTAL PLAIN¹

Lawn Grass Type	Monthly Nitrogen Application Rate per 1,000 Square Feet ²												Total lb N/ 1,000 sq ft/yr
	Jan	Feb	March	April	May	June	July	Aug ³	Sept	Oct	Nov	Dec	
BERMUDAGRASS													
Basic				1		1		1					3
High				1	1	1	1	1	1				6
CENTIPEDEGRASS³													
Basic					1								1
High					1			1					2
FESCUE, TALL													
Basic		0.5							1		0.5		2
High		1	0.5						1	1	0.5		4
ST. AUGUSTINEGRASS													
Basic				0.5	1	0.5	1	1	0.5	0.5			2
High				0.5	1	0.5	1	0.5	0.5				4
ZOYSIAGRASS (Emerald and Meyer cultivars)													
Basic				1	1	1		1					2
High				1	1	1		1					3
ZOYSIAGRASS (other cultivars)													
Basic				1	1	1		1					2
High				1	0.5	1	0.5	1					4

¹ All rates are per 1,000 sq ft. Multiply by 43.5 to convert to an acre basis. Follow table suggestions in the absence of soil test recommendations to the contrary. With the exception of centipedegrass, use a complete (N-P-K) turf-grade fertilizer in which 1/4 to 1/2 of the nitrogen is slowly available and that has a 3-1-2 or 4-1-2 analysis (e.g., 12-4-8, 16-4-8).

² In the absence of soil test recommendations, apply about 1 lb potassium per 1,000 sq ft using 1.6 lb of muriate of potash (0-0-60), 5 lb of sul-po-mag (0-0-22), or 2 lb of potassium sulfate (0-0-50) to bermudagrass, centipedegrass, and St. Augustinegrass.

³ Centipedegrass should be fertilized very lightly after establishment. Fertilize established centipedegrass using a low-phosphorus, high-potassium fertilizer with an analysis approaching 1-1-2 or 1-1-3. Fertilizers absent of phosphorus are preferred if soils supporting centipedegrass exhibit moderate to high levels of phosphorus.

TABLE 4-4B. SUGGESTED MAINTENANCE FERTILIZATION FOR CENTRAL PIEDMONT¹

Lawn Grass Type	Monthly Nitrogen Application Rate per 1,000 Square Feet ²												Total lb N/ 1,000 sq ft/yr
	Jan	Feb	March	April	May	June	July	Aug ³	Sept	Oct	Nov	Dec	
BERMUDAGRASS Basic High				1	1	1	1	1	1				3 6
CENTIPEDEGRASS³ Basic High					1			1					1 2
FESCUE, TALL Basic High		0.5 1	0.5						1 1	1	0.5 0.5		2 4
KENTUCKY BLUEGRASS Basic High		0.5 1	0.5						1 1	1	0.5 0.5		2 4
KENTUCKY BLUEGRASS- FINE FESCUE MIX Basic High		0.5 1	0.5						1 1	1	0.5 0.5		2 4
KENTUCKY BLUEGRASS- TALL FESCUE MIX Basic High		0.5 1	0.5						1 1	1	0.5 0.5		2 4
KENTUCKY BLUEGRASS- TALL FESCUE-FINE FESCUE MIX Basic High		0.5 1	0.5						1 1	1	0.5 0.5		2 4
KENTUCKY BLUEGRASS- PERENNIAL RYEGRASS MIX Basic High		1 1	0.5						1 1	1	1 0.5		3 4
ST. AUGUSTINEGRASS Basic High					1 1		1		1				2 3
ZOYSIAGRASS (Emerald and Meyer cultivars) Basic High				1	1		1						1 2
ZOYSIAGRASS (other cultivars) Basic High				1	1	1	1	1					2 3

¹ All rates are per 1,000 sq ft. Multiply by 43.5 to convert to an acre basis. Follow table suggestions in the absence of soil test recommendations to the contrary. With the exception of centipedegrass, use a complete (N-P-K) turf-grade fertilizer in which 1/4 to 1/2 of the nitrogen is slowly available and that has a 3-1-2 or 4-1-2 analysis (e.g., 12-4-8, 16-4-8).

² In the absence of soil test recommendations, apply about 1 lb potassium per 1,000 sq ft using 1.6 lb of muriate of potash (0-0-60), 5 lb of sul-po-mag (0-0-22), or 2 lb of potassium sulfate (0-0-50) to bermudagrass, centipedegrass, St. Augustinegrass, and zoysiagrass.

³ Centipedegrass should be fertilized very lightly after establishment. Fertilize established centipedegrass using a low-phosphorus, high-potassium fertilizer with an analysis approaching 1-1-2 or 1-1-3. Fertilizers absent of phosphorus are preferred if soils supporting centipedegrass exhibit moderate to high levels of phosphorus.

TABLE 4-4C. SUGGESTED MAINTENANCE FERTILIZATION FOR THE MOUNTAINS¹

Lawn Grass Type	Monthly Nitrogen Application Rate per 1,000 Square Feet ²												Total lb N/ 1,000 sq ft/yr
	Jan	Feb	March	April	May	June	July	Aug	Sept	Oct	Nov	Dec	
BERMUDAGRASS Basic High					1 1	1	1 1	1					2 4
FESCUE, TALL Basic High			0.5 1					1 1		0.5 1			2 3
KENTUCKY BLUEGRASS Basic High			1 1					1 1					2 3
KENTUCKY BLUEGRASS-FINE FESCUE MIX Basic High			1 1					1 1		1			2 3
KENTUCKY BLUEGRASS-TALL FESCUE MIX Basic High			1 1					1 1		1			2 3
KENTUCKY BLUEGRASS-TALL FESCUE-FINE FESCUE MIX Basic High			1 1					1 1		1			2 3
KENTUCKY BLUEGRASS- PERENNIAL RYEGRASS MIX Basic High			1 1					1 1	1	0.5 0.5			2.5 3.5
ZOYSIAGRASS (other cultivars) Basic High					0.5 1		1	0.5					1 2

¹ All rates are per 1,000 sq ft. Multiply by 43.5 to convert to an acre basis. Follow table suggestions in the absence of soil test recommendations to the contrary. With the exception of centipedegrass, use a complete (N-P-K) turf-grade fertilizer in which 1/4 to 1/2 of the nitrogen is slowly available and that has a 3-1-2 or 4-1-2 analysis (e.g., 12-4-8, 16-4-8).

² In the absence of soil test recommendations, apply about 1 lb potassium per 1,000 sq ft using 1.6 lb of muriate of potash (0-0-60), 5 lb of sul-po-mag (0-0-22), or 2 lb of potassium sulfate (0-0-50) to bermudagrass, St. Augustinegrass, and zoysiagrass.

Fertilizer Suggestions for Ornamental Plants in Landscapes

B. FAIR, L. BRADLEY and B. WHIPKER, Horticultural Science

The fertilizer suggestions given in the table are intended as a general guide. They are not a replacement for soil analyses from samples taken on a regular basis to assure maintenance of good nutrient availability and soil pH.

TABLE 4-5. FERTILIZER SUGGESTIONS FOR ORNAMENTAL PLANTS

Kind of Plant	Ratio	Analysis	Amount to Use	When to apply	Remarks
TREES*	3-1-2	12-4-8	1 to 2 lb N/inch trunk diameter	Apply fertilizer as a surface application to trees with nonrestricted root zone areas. Alleviate compaction of soil before applying. Another method is to drill holes 12 in. deep and 3 ft on center under tree. Place fertilizer and porous media into backfill. Water well. Best time to fertilize trees is in the fall.	Refer to Horticulture Information sheet 618, <i>Fertilizing Deciduous Shade Trees in the Landscape</i> for more information.
	or 1-1-1	8-8-8 or 10-10-10	1 to 2 lb N/1,000 sq ft of root zone		
SHRUBS	3-1-2	12-4-8	1 to 2 lb N/1,000 sq ft of bed area	Field grade fertilizers: Apply half in spring and the other half in summer.	Broadcast underneath shrubs in bed area and irrigate thoroughly after application. Keep fertilizer off foliage. It is not necessary to remove mulch if the area is irrigated properly.
	or 1-1-1	8-8-8 or 10-10-10	1 to 1.5 lb of complete fertilizer split in 2 applications per specimen shrub	Slow-release fertilizers: Apply entire amount as a single application before bud swell in the spring.	
FLOWERS	1-2-2	5-10-10	20 to 30 lb/1,000 sq ft for all analyses	Incorporate half the amount before planting; apply second half 6 weeks after planting.	Refer to Horticultural Information Leaflet 551, <i>Bed Preparation and Fertilizer Recommendations for Bedding Plants in the Landscape</i> .
	or 1-2-1	5-10-5			
	or 1-1-1	10-10-10			

* An alternate method is to use the appropriate complete fertilizer suggested above at planting time and each spring about the time growth starts. Then, if additional fertilizer is needed, use nitrogen. Example: per 1,000 sq ft, use approximately 6 lb of ammonium sulfate or 5 lb of ammonium nitrate.

Fertilizer Suggestions for Nursery Crops

T. E. BILDERBACK and A. V. LeBUDE, Horticultural Science; and R. GEHL, Soil Science

TABLE 4-6. FERTILIZER SUGGESTIONS FOR NURSERY CROPS

Kind of Production	Amount to Use ¹	Remarks
DECIDUOUS TREE SEEDLING BEDS	Year 1: 50 lb nitrogen (N) per acre or 18 oz N per 1,000 sq ft of bed. Year 2: Beds receive 100lb N per acre.	Year 1 : Surface apply after first true leaves appear. Base applications of other elements upon a soil test.
FIELD PRODUCTION	Year 1: 50 lb of nitrogen per acre incorporated as preplant. Year 2: 0.5 to 1 oz N per plant. Year 3: 1 to 2 oz N per plant. Do not exceed 100 to 200 lb N per acre. For liquid fertilizer applications through drip irrigation, reduce N application by 1/2 rate of annual field grade fertilizer. Apply equal rates of N during several irrigation events.	Year 1: Incorporate nutrients into the soil before planting. For field grade fertilizers, apply 0.67 of total amount before bud swell, 0.33 in June. Base applications of other elements upon a soil test. For more information, see http://www.ces.ncsu.edu/depts/hort/nursery/cultural/cultural_docs/field-bmps/building_nursery_soils.pdf
CONTAINER PRODUCTION	All essential elements must be provided when soilless mixes are used to produce nursery crops in containers. Controlled release fertilizers offer a consistent and reliable source of nutrients through the growing season. Formulations of nutrients and their release over time differ by fertilizer company and the optimal product chosen depends upon the species being grown as well as the location of the nursery in the state. Irrigation water quality can contribute significant amounts of some nutrients, such as calcium, magnesium, and iron. Test water quality of irrigation supplies at least once a year (e.g., mid-summer) to determine if nutrient adjustments are required. Diagnosis of whole production system nutritional problems requires analysis of foliar nutrient content (sample uppermost fully expanded leaves), collection of approximately 8 oz of leachate solution (collect from containers approximately 30 minutes after irrigation), and an irrigation water sample (collect from the irrigation head). Current research suggests that dolomitic limestone rates depend upon the calcium and magnesium content in irrigation water. As a result, limestone may not be required if provided by irrigation, however if dolomitic limestone is needed add only 2 to 5 lb per cubic yard of pine bark and sand potting mixes. Minor element supplements included in either NPK controlled release fertilizer products or from separate minor element packages are necessary in all pine bark mixes and should be incorporated if possible. Potting mixes containing composts, however, generally do not require dolomitic limestone or minor element supplements. Leachates, irrigation water, and foliar samples can be analyzed for \$5 each by the Agronomic Division, NCDA&CS, 4300 Reedy Creek Road, Raleigh, NC 27607-6465.	

¹ Rates may change with irrigation and soil type.

Lime and Fertilizer Suggestions for Vegetable Crops

J. R. SCHULTHEIS, and J. M. DAVIS, Horticultural Science;
C. R. CROZIER, R. GEHL, G. D. HOYT, and D. L. OSMOND, Soil Science

Important Notes:

1. For most vegetables grown on light-textured soils, apply the total recommended P₂O₅ and K₂O together with 25 to 50 percent of the recommended nitrogen before planting. The remaining nitrogen can be sidedressed with a fertilizer containing nitrogen only. Sidedressing or topdressing potash (K₂O) is recommended only on extremely light sandy soils with very low cation exchange capacities.
2. It may be desirable to build up the phosphorus and potassium levels in infertile loam and silt loam soils more rapidly than provided by these recommendations. In such instances, add an additional 40 to 50 pounds of P₂O₅ and K₂O, respectively, to the recommendations listed in the table for soils testing low in phosphorus and potassium. Apply the additional amounts as a broadcast and plow down or broadcast and disk-in application.
3. In the absence of soil tests, use recommendations listed under medium phosphorus and medium potassium levels on light-textured soils that have been in intensive vegetable production.
4. *For Piedmont growers producing vegetables on clay loam soils:* Reduce the recommended nitrogen and potassium rates by 20 percent and increase the phosphorus rate by 25 percent of the rates indicated in this table.

TABLE 4-7. LIME AND FERTILIZER SUGGESTIONS FOR VEGETABLE CROPS

Crop	Desirable pH	Nitrogen (N) lb/acre	Recommended Nutrients Based on Soil Tests								Total Amount of Nutrient Recommended and Suggested Methods of Application	
			Soil Phosphorus Level				Soil Potassium Level					
			Low	Med	High	Very High	Low	Med	High	Very High		
			P ₂ O ₅ lb/acre				K ₂ O lb/acre					
ASPARAGUS Growing crowns	6.5										Total recommended.	
		100	200	100	50	0	200	150	50	0	Broadcast and disk in.	
		50	200	100	50	0	100	75	50	0	Sidedress after cutting.	
		50	0	0	0	0	100	75	0	0	Sidedress after cutting.	
		New Planting Crowns and direct seeding	50	200	100	50	0	200	100	100	0	Total recommended.
			0	200	100	50	0	100	75	50	0	Broadcast and plow down.
			50	0	0	0	0	100	25	50	0	Sidedress at first cultivation.
		Cutting Bed or Nonhybrids	100	150	100	50	0	200	150	100	0	Total recommended.
			50	150	100	50	0	100	150	100	0	Broadcast and disk in.
			50	0	0	0	0	100	75	50	0	Sidedress at first cultivation.
		New hybrids	100	200	150	100	0	300	225	150	0	Total recommended.
			50	200	150	100	0	150	100	75	0	Broadcast before cutting season.
			50	0	0	0	0	100	125	75	0	Sidedress after cutting.
Apply 2 lb boron (B) per acre every 3 years on most soils.												
BEAN, Lima Single crop	6 to 6.5	70 to 110	120	80	40	20	160	120	80	20	Total recommended.	
		25 to 50	80	40	20	0	120	80	60	0	Broadcast and disk-in.	
		20	40	40	20	20	40	40	20	20	Band-place with planter.	
		25 to 40	0	0	0	0	0	0	0	0	Sidedress 3 to 5 weeks after emergence.	
BEAN, Snap	6 to 6.5	40 to 80	80	60	40	20	80	60	40	20	Total recommended.	
		20 to 40	40	40	0	0	40	40	0	0	Broadcast and disk-in.	
		20 to 40	40	20	40	20	40	20	40	20	Band-place with planter.	
BEET	6 to 6.5	75 to 100	150	100	50	0	150	100	50	0	Total recommended.	
		50	150	100	50	0	150	100	50	0	Broadcast and disk-in.	
		25 to 50	0	0	0	0	0	0	0	0	Sidedress 4 to 6 weeks after planting.	
BROCCOLI	6 to 6.5	125 to 175	200	100	50	0	200	100	50	0	Total recommended.	
		50 to 100	150	100	50	0	150	100	50	0	Broadcast and disk-in.	
		50	50	0	0	0	50	0	0	0	Sidedress 2 to 3 weeks after planting.	
		25	0	0	0	0	0	0	0	0	Sidedress every 2 to 3 weeks after first sidedressing.	
Apply 2 lb boron (B) per acre with broadcast fertilizer.												
BRUSSEL SPROUT, CABBAGE, and CAULIFLOWER	6 to 6.5	100 to 175	200	100	50	0	200	100	50	0	Total recommended.	
		50 to 75	200	100	50	0	200	100	50	0	Broadcast and disk-in.	
		25 to 50	0	0	0	0	0	0	0	0	Sidedress 2 to 3 weeks after planting.	
		25 to 50	0	0	0	0	0	0	0	0	Sidedress if needed, according to weather.	
Apply 2 to 3 lb boron (B) per acre and molybdenum (mo) per acre as 0.5 lb sodium molybdate per acre with broadcast fertilizer.												

TABLE 4-7. LIME AND FERTILIZER SUGGESTIONS FOR VEGETABLE CROPS

Crop	Desirable pH	Nitrogen (N) lb/acre	Recommended Nutrients Based on Soil Tests								Total Amount of Nutrient Recommended and Suggested Methods of Application
			Soil Phosphorus Level				Soil Potassium Level				
			Low	Med	High	Very High	Low	Med	High	Very High	
			P ₂ O ₅ lb/acre				K ₂ O lb/acre				
CARROT	6 to 6.5	50 to 80	150	100	50	0	150	100	50	0	Total recommended.
		50	150	100	50	0	150	100	50	0	Broadcast and disk-in.
		25 to 30	0	0	0	0	0	0	0	0	Sidedress if needed.
		Apply 1 to 2 lb boron (B) per acre with broadcast fertilizer.									
CELERY	6 to 6.5	75 to 100	250	150	100	0	250	150	100	0	Total recommended.
		50	250	150	100	0	250	150	100	0	Broadcast and disk-in or drill deep.
		25 to 50	0	0	0	0	0	0	0	0	Sidedress 2 to 3 weeks after planting.
		Apply 2 to 3 lb boron (B) per acre with broadcast fertilizer.									
CORN, Sweet	6 to 6.5	110 to 155	160	120	80	20	160	120	80	20	Total recommended.
		40 to 60	120	100	60	0	120	100	60	0	Broadcast before planting.
		20	40	20	20	20	40	20	20	20	Band-place with planter.
		50 to 75	0	0	0	0	0	0	0	0	Sidedress when corn is 12 to 18 in. tall.
		Apply 1 to 2 lb boron (B) per acre with broadcast fertilizer. NOTE: On very light sandy soils, sidedress 40 lb N per acre when corn is 6 in. tall and another 40 lb N per acre when corn is 12 to 18 in. tall.									
CUCUMBER	6 to 6.5	80 to 160	150	100	50	25	200	150	100	25	Total recommended.
		40 to 100	125	75	25	0	175	125	75	0	Broadcast and disk-in.
		20 to 30	25	25	25	25	25	25	25	25	Band-place with planter 7 to 14 days after planting.
		20 to 30	0	0	0	0	0	0	0	0	Sidedress when vines begin to run, or apply in irrigation water.
		Drip fertilization: See "cucumber" in specific recommendations in the 2012 Southeastern Vegetable Crop Handbook.									
EGGPLANT Bareground Plasticulture	6 to 6.5	100 to 200	250	150	100	0	250	150	100	0	Total recommended.
		50 to 100	250	150	100	0	250	150	100	0	Broadcast and disk-in.
		25 to 50	0	0	0	0	0	0	0	0	Sidedress 3 to 4 weeks after planting.
		25 to 50	0	0	0	0	0	0	0	0	Sidedress 6 to 8 weeks after planting.
	Apply 1 to 2 lb boron (B) per acre with broadcast fertilizer..										
	Plasticulture	145	250	150	100	0	240	170	100	0	Total recommended.
		50	250	150	100	0	100	100	100	0	Broadcast and disk in.
		95	0	0	0	0	140	70	0	0	Fertigate.
	Apply 1 to 2 lb boron (B) per acre with broadcast fertilizer. Drip fertilization: See "eggplant" in specific recommendations in the 2012 Southeastern Vegetable Crop Handbook.										
	ENDIVE, ESCAROLE, LEAF LETTUCE	6 to 6.5	75 to 125	200	150	100	0	200	150	100	0
50 to 75			200	150	100	0	200	150	100	0	Broadcast and disk-in.
25 to 50			0	0	0	0	0	0	0	0	Sidedress 3 to 5 weeks after planting.
ICEBERG LETTUCE	6 to 6.5	85 to 175	200	150	100	0	200	150	100	0	Total recommended.
		60 to 80	200	150	100	0	200	150	100	0	Broadcast and disk-in.
		25 to 30	0	0	0	0	0	0	0	0	Sidedress 3 times beginning 2 weeks after planting.
LEAFY GREENS, COLLARD, KALE, MUSTARD	6 to 6.5	75 to 80	150	100	50	0	150	100	50	0	Total recommended.
		50	150	100	50	0	150	100	50	0	Broadcast and disk-in.
		25 to 30	0	0	0	0	0	0	0	0	Sidedress, if needed.
		Apply 1 to 2 lb boron (B) per acre with broadcast fertilizer.									
LEEK	6 to 6.5	75 to 125	200	150	100	0	200	150	100	0	Total recommended.
		50 to 75	200	150	100	0	200	150	100	0	Broadcast and disk-in.
		25 to 50	0	0	0	0	0	0	0	0	Sidedress 3 to 4 weeks after planting if needed.
		Apply 1 to 2 lb boron (B) per acre with broadcast fertilizer.									
CANTALoupES AND MIXED MELONS Bareground	6 to 6.5	75 to 115	150	100	50	25	200	150	100	25	Total recommended.
		25 to 50	125	75	25	0	175	125	75	0	Broadcast and disk-in.
		25	25	25	25	25	25	25	25	25	Band-place with planter.
		25 to 40	0	0	0	0	0	0	0	0	Sidedress when vines start to run.
		Apply 1 to 2 lb boron (B) per acre with broadcast fertilizer.									

TABLE 4-7. LIME AND FERTILIZER SUGGESTIONS FOR VEGETABLE CROPS

Crop	Desirable pH	Nitrogen (N) lb/acre	Recommended Nutrients Based on Soil Tests								Total Amount of Nutrient Recommended and Suggested Methods of Application
			Soil Phosphorus Level				Soil Potassium Level				
			Low	Med	High	Very High	Low	Med	High	Very High	
			P ₂ O ₅ lb/acre				K ₂ O lb/acre				
CANTALOUPE AND MIXED MELONS Plasticulture	6 to 6.5	75 to 150	150	100	50	25	200	150	100	25	Total recommended.
		25	150	100	50	25	100	75	50	25	Broadcast and disk in.
		50 to 100	0	0	0	0	100	75	50	0	Fertigate.
		Apply 1 to 2 lb boron (B) per acre with broadcast fertilizer. Drip fertilization: See "muskmelon" in specific recommendations in the 2012 Southeastern Vegetable Crop Handbook.									
OKRA	6 to 6.5	100 to 200	250	150	100	0	250	150	100	0	Total recommended.
		50 to 100	250	150	100	0	250	150	100	0	Broadcast and disk-in.
		25 to 50	0	0	0	0	0	0	0	0	Sidedress 3 to 4 weeks after planting.
		25 to 50	0	0	0	0	0	0	0	0	Sidedress 6 to 8 weeks after planting.
	Apply 1 to 2 lb boron (B) per acre with broadcast fertilizer. NOTE: Where plastic mulches are being used, broadcast 50 to 100 lb nitrogen (N) per acre with recommended P ₂ O ₅ and K ₂ O and disk incorporate prior to laying mulch. Drip fertilization: See "okra" in specific recommendations in the 2012 Southeastern Vegetable Crop Handbook.										
ONION Bulb Green	6 to 6.5	75 to 125	200	100	50	0	200	100	50	0	Total recommended.
		50 to 75	200	100	50	0	200	100	50	0	Broadcast and disk-in.
		25 to 50	0	0	0	0	0	0	0	0	Sidedress 4 to 5 weeks after planting.
		150 to 175	200	100	50	0	200	100	50	0	Total recommended.
		50 to 75	200	100	50	0	200	100	50	0	Broadcast and disk-in.
		50	0	0	0	0	0	0	0	0	Sidedress 4 to 5 weeks after planting.
		50	0	0	0	0	0	0	0	0	Sidedress 3 to 4 weeks before harvest.
	Apply 1 to 2 lb boron (B) and 20 lb sulfur (S) per acre with broadcast fertilizer.										
PARSLEY	6 to 6.5	100 to 175	200	150	100	0	200	150	100	0	Total recommended.
		50 to 75	200	150	100	0	200	150	100	0	Broadcast and disk-in.
		25 to 50	0	0	0	0	0	0	0	0	Sidedress after first cutting.
		25 to 50	0	0	0	0	0	0	0	0	Sidedress after each additional cutting.
PARSNIP	6 to 6.5	50 to 100	150	100	50	0	150	100	50	0	Total recommended.
		25 to 50	150	100	50	0	150	100	50	0	Broadcast and disk-in.
		25 to 50	0	0	0	0	0	0	0	0	Sidedress 4 to 5 weeks after planting.
			Apply 1 to 2 lb boron (B) per acre with broadcast fertilizer.								
PEA, English Spring plowed	5.8 to 6.5	40 to 60	120	80	40	0	120	80	40	0	Total recommended. Broadcast and disk-in before seeding.
PEA, Southern	5.8 to 6.5	16	96	48	0	0	96	48	0	0.	Broadcast and disk-in.
PEPPER Bareground Plasticulture	6 to 6.5	100 to 130	200	150	100	0	200	150	100	0	Total recommended.
		50	200	150	100	0	200	150	100	0	Broadcast and disk-in.
		25 to 50	0	0	0	0	0	0	0	0	Sidedress after first fruit set.
		25 to 30	0	0	0	0	0	0	0	0	Sidedress later in season if needed.
		100 to 185	200	150	100	0	365	300	235	0	Total recommended
		50	200	150	100	0	100	100	100	0	Broadcast and disk in.
		50 to 135	0	0	0	0	265	200	135	0	Fertigate
	Drip fertilization: See "pepper" in specific commodity recommendations in <i>Plasticulture for Commercial Vegetables</i> (AG-489).										
POTATO, Irish Loams and silt loams Sandy loams and loamy sands	5.8 to 6.2	100 to 150	110	90	70	50	200	150	50	50	Total recommended.
		85 to 135	60	40	20	0	200	150	50	50	Broadcast and disk in.
		15	50	50	50	50	0	0	0	0	Band-place with planter at planting.
		150	200	150	100	50	300	200	100	50	Total recommended.
		50	200	150	100	50	300	200	100	50	Broadcast and disk in.
PUMPKIN and SQUASH (Winter) Bareground Plasticulture	6 to 6.5	80 to 90	150	100	50	0	200	150	100	0	Total recommended.
		40 to 50	150	100	50	0	200	150	100	0	Broadcast and disk in.
		40 to 45	0	0	0	0	0	0	0	0	Sidedress when vines begin to run.
		80 to 150	150	100	50	0	200	150	100	0	Total recommended.
		25 to 50	150	100	50	0	100	75	50	0	Disk in row
RADISH	6 to 6.5	55 to 100	0	0	0	0	100	75	50	0	Sidedress when vines begin to run.
		50	150	100	50	0	150	100	50	0	Total recommended. Broadcast and disk-in or drill deep.
	Apply 1 to 2 lb boron (B) per acre with broadcast fertilizer.										

TABLE 4-7. LIME AND FERTILIZER SUGGESTIONS FOR VEGETABLE CROPS

Crop	Desirable pH	Nitrogen (N) lb/acre	Recommended Nutrients Based on Soil Tests								Total Amount of Nutrient Recommended and Suggested Methods of Application
			Soil Phosphorus Level				Soil Potassium Level				
			Low	Med	High	Very High	Low	Med	High	Very High	
			P ₂ O ₅ lb/acre				K ₂ O lb/acre				
RUTABAGA and TURNIP	6 to 6.5	50 to 75	150	100	50	0	150	100	50	0	Total recommended.
		25 to 50	150	100	50	0	150	100	50	0	Broadcast and disk in.
		25 to 50	0	0	0	0	0	0	0	0	Sidedress when plants are 4 to 6 in. tall.
		Apply 1 to 2 lb boron (B) per acre with broadcast fertilizer.									
SPINACH Fall Overwinter	6 to 6.5	75 to 125	200	150	100	0	200	150	100	0	Total recommended.
		50 to 75	200	150	100	0	200	150	100	0	Broadcast and disk in.
		25 to 50	0	0	0	0	0	0	0	0	Sidedress or topdress.
		80 to 120	0	0	0	0	0	0	0	0	Total recommended for spring application on overwintered crop.
		50 to 80	0	0	0	0	0	0	0	0	Apply in late February.
		30 to 40	0	0	0	0	0	0	0	0	Apply in late March.
SQUASH, Summer	6 to 6.5	100 to 130	150	100	50	0	200	150	100	0	Total recommended.
		25 to 50	150	100	50	0	150	100	50	0	Broadcast and disk in.
		50	0	0	0	0	0	0	0	0	Sidedress when vines start to run.
		25 to 30	0	0	0	0	0	0	0	0	Apply through irrigation system.
Apply 1 to 2 lb boron (B) per acre with broadcast fertilizer. Drip/trickle fertilization: See "summer squash" in specific recommendations in the 2012 Southeastern Vegetable Crop Handbook.											
SWEETPOTATO	5.8 to 6.2	50 to 80	200	100	50	0	300	200	150	0	Total recommended.
		0	150	60	30	0	150	50	30	0	Broadcast and disk in.
		50 to 80	0	0	0	0	150	150	120	120	Sidedress 21 to 28 days after planting.
		Add 0.5 lb of actual boron (B) per acre 40 to 80 days after planting.									
TOMATO Bare ground for sandy loams and loamy sands Bareground for loams and clay	6 to 6.5	80 to 90	200	150	100	0	300	200	100	0	Total recommended.
		40 to 45	200	150	100	0	300	200	100	0	Broadcast and disk in.
		40 to 45	0	0	0	0	0	0	0	0	Sidedress when first fruits are set and as needed.
	6 to 6.5	75 to 80	200	150	100	0	250	150	100	0	Total recommended.
		50	200	150	100	0	250	150	100	0	Broadcast and plow down.
		25 to 30	0	0	0	0	0	0	0	0	Sidedress when first fruits are set and as needed.
Apply 1 to 2 lb boron (B) per acre with broadcast fertilizer.											
TOMATO Plasticulture		130 to 210	200	150	100	0	420	325	275	0	Total recommended.
		50	200	150	100	0	295	220	125	0	Broadcast and disk in.
		80 to 160	0	0	0	0	295	220	125	0	Fertigate
Apply 1 to 2 lb boron (B) per acre with broadcast fertilizer. Drip fertilization: See "tomato" in specific recommendations in the 2012 Southeastern Vegetable Crop Handbook.											
WATERMELON Nonirrigated Irrigated Plasticulture	6 to 6.5	75 to 90	150	100	50	0	200	150	100	0	Total recommended.
		50	150	100	50	0	200	150	100	0	Broadcast and disk in.
		25 to 40	0	0	0	0	0	0	0	0	Topdress when vines start to run.
		100 to 150	150	100	50	0	200	150	100	0	Total recommended.
		50	150	100	50	0	150	150	100	0	Broadcast and disk in.
		25 to 50	0	0	0	0	0	0	0	0	Topdress when vines start to run.
		25 to 50	0	0	0	0	0	0	0	0	Topdress at first fruit set.
		125 to 150	150	100	50	0	200	150	100	0	Total recommended.
		25 to 50	150	100	50	0	100	75	50	0	Disk in row.
		100	0	0	0	0	100	75	50	0	Fertigation
NOTE: Excessive rates of N may increase hollow heart in seedless watermelons. Drip fertilization: See "watermelon" in specific recommendations in the 2012 Southeastern Vegetable Crop Handbook.											

Fertilizer Rules and Regulations

D. L. OSMOND, C. R. CROZIER, R. GEHL, and D.A. CROUSE, Soil Science Department; D. H. HARDY, NCDA&CS

Fertilizer, lime and landplaster are regulated by Plant Industry- NCDA&CS and rules and regulations in their entirety are found at <http://www.ncagr.gov/plantindustry/pubs.htm>.

Chlorine Guarantees for Tobacco Fertilizer

The maximum chlorine (Cl) guarantees permitted for tobacco plantbed fertilizer shall be:

1. For fertilizers with a nitrogen (N) guarantee up to and including 6 percent, 0.5 percent chlorine (Cl).
2. For fertilizers with a nitrogen (N) guarantee above 6 percent, 1 percent chlorine (Cl).

The maximum chlorine (Cl) guarantees permitted on field crop tobacco fertilizer shall be:

1. For fertilizer with a nitrogen (N) guarantee up to and including 4 percent, a maximum chlorine (Cl) guarantee of 2 percent.
2. For fertilizer with a nitrogen (N) guarantee greater than 4 percent, a maximum percent chlorine (Cl) guarantee not more than one-half of the respective total nitrogen (N) guarantee.

The maximum chlorine (Cl) permitted in tobacco top-dressers shall be 2 percent.

Size Standards for Agricultural Liming Material

Agricultural liming material shall conform to the following minimum screening standards:

1. Ninety percent must pass through U.S. Standard 20-mesh screen, with a tolerance of 5 percent.
2. For dolomitic limestone, 35 percent must pass through U.S. Standard 100-mesh screen; for calcitic limestone, 25 percent must pass through U.S. Standard 100-mesh screen, with a tolerance of 5 percent.

Additional Criteria for Agricultural Liming Material

1. A product must contain at least 6 percent magnesium (Mg) from magnesium carbonate to be classified as a dolomitic limestone.
2. There is no minimum calcium carbonate equivalent (CCE) requirement for limestone sold in North Carolina. However, the product must be labeled to show the amount necessary to equal that required from a liming material having a 90 percent CCE. Lime recommendations in North Carolina are based on 90 percent CCE. For example, a product having a CCE of 50 percent would be labeled "3,600 lb of this material equals 1 ton of standard agricultural liming material."
3. Pelleted lime must slake down when it comes in contact with moisture, thereby meeting the size standards for agricultural liming materials given above.

Nutrient Content of Fertilizer Materials

D. L. OSMOND, R. J. GEHL, and C. R. CROZIER, Soil Science; T. E. BILDERBACK, Horticultural Science

Although fertilizer materials may contain two or more nutrients, usually only one of the nutrients is commonly associated with the material. The content of that nutrient is generally well-known, but the contents of other associated nutrients are less well-known. Information on the contents of lesser-known nutrients in fertilizer materials exists but is often not readily available. Information presented below gives the nutrients and their contents that are normally found in fertilizer materials. In the case of micronutrient fertilizers, only the principle micronutrient is considered relevant. This information should be considered only as a guide, since actual nutrient contents may vary slightly, depending on the source of the fertilizer material. Additional information about the nutrient content of fertilizer and organic materials can be found in the N.C. Cooperative Extension Service publications AG-439-18, AG-439-19, and AG-467 or on the Internet at http://www.bae.ncsu.edu/bae/programs/extension/publicat/wqwm/ag473_14.html.

TABLE 4-8. COMPOSITION OF SELECTED MICRONUTRIENT FERTILIZERS

Element	Fertilizer	Chemical Formula	Elemental Percentage (%)
Mn	Manganese sulfate	MnSO ₄ •3H ₂ O	27
	Manganous oxide ²	MnO	41 to 68
	Manganese chloride	MnCl ₂	7
	Manganese oxide ²	MnO ₂	62 to 70
Mo	Sodium molybdate	Na ₂ MoO ₄ •H ₂ O	38
Cu	Copper sulfate	CuSO ₄ •5H ₂ O	25
Zn	Zinc sulfate	ZnSO ₄ •H ₂ O	22 to 36
	Zinc oxide ²	ZnO	78
Fe	Ferrous sulfate	FeSO ₄ •7H ₂ O	20
	Ferric sulfate	Fe ₂ (SO ₄) ₃ •4H ₂ O	23
B	Borax	Na ₂ B ₄ O ₇ •10H ₂ O	11
	Solubor	Na ₂ B ₈ O ₁₃ •4H ₂ O	20

TABLE 4-9. COMPOSITION OF SELECTED ORGANIC FERTILIZER MATERIALS

Nutrient Source	Analysis (%) ¹			Relative Rate of Nutrient Release
	N	P ₂ O ₅	K ₂ O	
Animal tankage	7	9	0	Medium
Bone meal				
Raw	3	22	0	Very slow
steamed	2	28	0	Slow
Castor pomace	5	1	1	Slow
Cotton seed meal	7	2	2	Slow
Dried blood	12	3	0	Fast
Feather meal	14	0.3	0.1	Fast
Hardwood ashes ₂	0.5	1	5	Slow
Linseed meal	5	2	2	Slow
Municipal yard and leaf compost ⁴	0.5	0.4	0.8	Slow
Sheep wool pellets	5.8	0.9	2.4	Fast/Slow
Softwood ashes ²	0.5	2	4	Slow
Tobacco stems	2	1	7	Slow
Turkey litter compost ³	1.5	3.5	1	Slow

¹ Percentages of nutrient elements may vary depending upon source. Values given are approximations.

² Plant-available N less than or equal to 0.25%.

³ Compost products must be stabilized before use, or they may cause nutrient depletion.

⁴ Municipal yard and leaf waste composts may contain measurable concentrations of metals, such as lead and zinc. A complete analysis of these soil amendments should be conducted before use.

TABLE 4-10. COMPOSITION OF TYPICAL N SOLUTIONS¹

Contents (%)	No-Pressure Nonammonia Solutions ²					Low Pressure Ammonia Solution	Aqua-Ammonia
	16.0	19.0	21.0	30.0	32.0	37.0	20.0
Total N	16.0	19.0	21.0	30.0	32.0	37.0	20.0
NH ₃	0	0	0	0	0	16.6	24.4
NH ₄ NO ₃	45.8	54.3	60.0	42.2	44.3	66.8	0
Urea	0	0	0	32.7	35.4	0	0
Water	54.2	45.7	40.0	25.1	20.3	16.6	75.6
NO ₃ -N	8.0	9.5	10.5	7.4	7.7	11.7	0
NH ₄ -N	8.0	9.5	10.5	7.4	7.8	25.4 ³	20.0 ³
Urea-N	0	0	0	15.2	16.4	0	0
	Salt-out Temperatures (F) ⁴						
		41	41	15	32	56	-58

¹ Values may vary depending on product.

² Several companies provide N solutions containing 3.5% to 5.0% sulfur.

³ Includes N present as NH₃.

⁴ Proprietary additives can alter salt-out temperatures; this should be verified with the dealer.

TABLE 4-11. COMPOSITION OF SELECTED FERTILIZER MATERIALS¹

Fertilizer Material	Chemical Formula	Nutrient Percentage (%)							
		NO ₃ -N	NH ₄ -N	Total N	P ₂ O ₅	K ₂ O	Ca ⁴	Mg	S
Ammonium nitrate	NH ₄ NO ₃	17	17	34					
Monoammonium phosphate	NH ₄ H ₂ PO ₄		11	11	48		1		2
Diammonium phosphate	(NH ₄) ₂ HPO ₄		16 to 18	16 to 18	46 to 48				
Ammonium sulfate	(NH ₄) ₂ SO ₄		21	21					24
Ammonium thiosulfate	(NH ₄) ₂ S ₂ O ₃			12					26
Anhydrous ammonia	NH ₃		82	82					
Urea-Form ²			38	38					
Calcium nitrate	Ca(NO ₃) ₂	15		15			19	1	
Nitrate of soda potash	NaNO ₃ •KNO ₃	15		15		14			
Sodium nitrate	NaNO ₃	16		16					
Urea	CO(NH ₂) ₂ •H ₂ O		45 to 46	45 to 46					
Single superphosphate	Ca(H ₂ PO ₄) ₂ + CaSO ₄				18 to 20		18 to 21		12
Triple superphosphate	Ca(H ₂ PO ₄) ₂ •H ₂ O				42 to 50		12 to 14		1
Basic slag ³	5CaO•P ₂ O ₅ •SiO				2 to 17		3 to 33	3	
Potassium chloride	KCl					60 to 62			
Potassium nitrate	KNO ₃	13		13		44			
Potassium sulfate	K ₂ SO ₄					50 to 53		1	18
Potassium-magnesium sulfate	K ₂ SO ₄ •2MgSO ₄					22		11	23
Epsom salt	MgSO ₄ •7H ₂ O						2	10	14
Gypsum	CaSO ₄ •2H ₂ O						23		18.5

¹ Values may vary depending on product.

² Slow release N source.

³ Lime value — about 0.67 agricultural limestone.

⁴ Evaluate gypsum products marked as Ca sources based on guaranteed fertilizer analysis. Pure calcium sulfate is 29 percent calcium. Landplasters typically contain 70 to 85 percent calcium sulfate (21 to 25 percent calcium), while phosphogypsum and other products may contain 50 percent or less calcium sulfate (15 percent calcium). See Table 4-1 for rate recommendations for peanut.

Solubility of Selected Fertilizer Materials

To be available to plants, at least some of a nutrient must be slightly soluble in the soil solution. The amount of substance that will dissolve at a given temperature in water is known as its solubility. Solubility of most chemicals is slightly higher at higher temperatures; that of others, especially ammonium and potassium nitrates, increases rapidly with temperature. The presence of other substances in the solution may either increase or decrease solubility. The solubility of selected pure fertilizer materials in water at 32°F is shown below.

TABLE 4-12. SOLUBILITY OF SELECTED FERTILIZER MATERIALS

Fertilizer Material	Chemical Formula	Solubility (lb/100 gal)	Salt Index (relative effect on the soil solution)
Ammonia	NH ₃	750	47.1
Ammonium nitrate	NH ₄ NO ₃	983	104.7
Ammonium sulfate	(NH ₄) ₂ SO ₄	592	69.0
Borax	Na ₂ B ₄ O ₇ •10H ₂ O	25	
Calcium carbonate (limestone)	CaCO ₃	0.050	4.7
Calcium metaphosphate	Ca(PO ₃) ₂	0.008	
Calcium nitrate	Ca(NO ₃) ₂ •4H ₂ O	1,117	52.5
Calcium sulfate	CaSO ₄ •2H ₂ O	2	8.1
Copper sulfate	CuSO ₄ •5H ₂ O	267	
Diammonium phosphate	(NH ₄) ₂ HPO ₄	209	29.9
Dicalcium phosphate	CaHPO ₄ •2H ₂ O	0.168	8
Magnesia	MgO	0.005	1.4
Magnesium sulfate	MgSO ₄ •7H ₂ O	580	44
Manganese sulfate	MnSO ₄ •4H ₂ O	875	
Monoammonium phosphate	NH ₄ H ₂ PO ₄	358	34.2
Monocalcium phosphate	CaH ₄ (PO ₄) ₂ •H ₂ O	*	15.4
Potassium chloride	KCl	233	116.3
Potassium-magnesium sulfate	K ₂ SO ₄ •2MgSO ₄	200	
Potassium nitrate	KNO ₃	108	73.6
Potassium sulfate	K ₂ SO ₄	67	46.1
Sodium nitrate	NaNO ₃	608	100.0
Urea	CO(NH ₂) ₂	559	75.4
Zinc sulfate	ZnSO ₄ •6H ₂ O	584	

* It decomposes with a small amount of water and is soluble in a large amount. The solubility varies with the conditions.

Mixing Herbicides with Nitrogen Solutions or Fluid Fertilizers

D. L. JORDAN, Crop Science; C. R. CROZIER and R. J. GEHL, Soil Science

Tank-mixing and applying herbicides with nonpressure nitrogen solutions or fluid fertilizers may offer savings in labor and time by eliminating at least one trip over the field. Effectiveness of some postemergence-directed herbicides for corn and sorghum is increased by applying them in nitrogen solution.

Herbicide labels will indicate if the product can be mixed with sprayable fertilizer. Herbicides may not always mix evenly throughout a sprayable fluid fertilizer, or the components may separate too quickly to make their combined use practical. Therefore, test every batch of fertilizer for compatibility before adding the herbicide. Batches can vary in pH, salt content, and salt concentration. Even these minor differences may affect compatibility. Sulfur-containing nitrogen solutions can be particularly troublesome.

The information presented below is intended as a guide only. The term "compatibility" as used here refers to chemical and physical compatibility and is not intended to supplant label directions.

A guide for making your own compatibility test follows. Also, herbicide labels give information on compatibility, tank-mix combinations, and procedures for testing compatibility with sprayable fluid fertilizer. Be sure to read the label!

Precautions

To help ensure successful results from application of herbicides and nonpressure nitrogen solutions or fluid fertilizers:

- Always check compatibility by making a small scale test (directions given below) before mixing in field.
- Use a compatibility agent if indicated on the herbicide label or if your small-scale test indicates the need.
- Vigorously agitate tank contents while mixing and applying. The spray application equipment should use a high-capacity (eight-roller, internal gear, or centrifugal) pump. Spray tanks should be equipped with hydraulic jet agitators mounted in the bottom of the tank. Bypass lines usually do not provide adequate agitation. Jet agitators should be attached to a separate line from the pump.
- Do not connect agitators to by-pass lines. For simple hydraulic jet agitators, a flow rate of 6 gallons per minute for every 100 gallons of tank capacity is sufficient. If volume-booster nozzles are used for agitation, the flow rate can be reduced to 2 to 3 gallons per minute for every 100 gallons of tank capacity. If a liquid-fertilizer type applicator is used with a metering pump, agitation must also be supplied. The best method is to use a separate power takeoff pump for extra circulation.
- If a wettable powder or dry flowable is used, make a slurry with water and add slowly to the tank. Add wettable powder first, dry flowables second, flowables third, and liquids last.
- A number of dry flowable herbicides now come packaged in water-soluble film packets. These packets usually will not dissolve in nitrogen solution or fluid fertilizer. When using nitrogen solution or fluid fertilizer as the carrier for a product packed in these film packets, slurry the herbicide in clean water before adding to the spray tank.
- If a flowable product is used, premix one part flowable with one part water and add diluted mixture slowly to tank. The fluid fertilizer may be substituted for the water after compatibility has been checked.
- Premix liquid products with two parts water or the fertilizer carrier before adding into the tank.
- Reduce drift by choosing an appropriate nozzle size to put out the desired amount of solution without causing excessive pressure.
- Caution:* Take extra care in applying herbicide-fluid fertilizer mixtures to ensure that the correct herbicide rate is applied, that the herbicide is distributed uniformly, and that all directions concerning application of the herbicide are followed. Sprayer output will not be the same using fluid fertilizer as when using water as the carrier. Recalibrate sprayers for the fertilizer carrier. Do not apply herbicide-fluid fertilizer mixes overtop any crop except small grains, as injury will result.

How to Test Compatibility of Herbicides with Fluid Fertilizers

D. L. JORDAN, Crop Science; C. R. CROZIER and R. J. GEHL, Soil Science

Follow the compatibility test procedures on the herbicide label. If not on the label, follow the directions listed below.

1. Put 1 pint of fluid fertilizer in each of two 1-quart jars.
2. Following the adjacent table, add 0.25 to 0.375 teaspoon of a compatibility agent to one jar and shake for 5 or 10 seconds to mix. One-fourth teaspoon is equivalent to 2 pt/100 gallons of fluid fertilizer. Mark the jar "with" to indicate the compatibility agent has been added.
3. Next, add the proper amount of herbicide to each jar, according to the table. For herbicides used in small quantities per acre, one will need a greater volume of fluid fertilizer for the compatibility test. Adjust the herbicide/fertilizer ratio as follows:

1 oz dry flowable per acre = 0.7 tsp in 1 gal

1 fl oz liquid per acre = 0.25 tsp in 1 gal

If more than one herbicide is to be used in the mixture, add them separately with the wettable powders or dry flowables first, flowables second, and liquids last. Shake

the jar gently for 5 to 10 seconds after each addition.

4. Let the jars stand for 5 minutes, then check for formation of large flakes, sludge, gels, or other precipitates, or to see if the herbicide remains as small, oily particles in the solution.
5. Allow both jars to stand for 30 minutes, checking them periodically. An emulsifiable concentrate normally will go to the top after standing, whereas wettable powders or dry flowables will either settle to the bottom of the jar or float to the top, depending on the density of the fertilizer and the herbicide. If the separate layers of fluid fertilizers and additives (herbicide and compatibility agent) can be resuspended by shaking, commercial application is possible.
6. If incompatibility of any form occurs in the jar with the compatibility agent added, do not mix the fluid fertilizer and herbicide in the same spray tank. If incompatibility occurs only in the jar without the compatibility agent, use of a compatibility agent is recommended.

TABLE 4-13. Amounts of Herbicides to Use to Test Compatibility¹

Application Rate Product Per Acre	Add to 1 Pint of Fluid Fertilizer
Wettable powders or dry herbicides	
1 lb	1.4 tsp
2 lb	2.9 tsp
3 lb	4.3 tsp
4 lb	5.8 tsp
5 lb	7.2 tsp
Emulsifiable concentrates, flowables, liquids, or solutions	
1 pt	0.5 tsp
1 qt	0.9 tsp
2 qt	1.9 tsp
3 qt	2.9 tsp
1 gal	3.8 tsp
5 qt	4.8 tsp

¹This compatibility test is designed for 25 gal of spray per acre with the maximum labeled rate of herbicide. For changes in spray volume or herbicide concentration, make the appropriate proportional change in the ingredients in the test. Regardless of spray volume, the amount of compatibility agents should be equal to 2 to 3 pt (0.25 tsp equals 2 pt; 0.375 tsp equals 3 pt) per 100 gal of fertilizer.

Fertilizer Placement

C. R. CROZIER, D. L. OSMOND, and R. J. GEHL, Soil Science, and D. H. HARDY, NCDA&CS

Proper fertilizer placement can be as important as what kind of fertilizer you use. Fertilizer placed in contact with or too close to seeds and young plants can cause salt injury, resulting in poor stands and slow starts. Nonuniform broadcasting of fertilizer with shallow mixing just before planting may give streaks through the field due to delayed germination or seedling injury. Salt injury is most severe in dry weather or following light rains that dissolve the fertilizer salts and leave highly concentrated salt solutions in the root zone. Nitrogen and potassium salts account for most of this injury.

To reduce salt injury risk, a side-band placement of starter fertilizer at planting is generally preferred over application directly in contact with the seed, commonly referred to as "pop-up" placement. Good results are often achieved for seeded crops like corn, cotton, sorghum, and soybeans using a 2 by 2-inch placement, placing the fertilizer 2 inches to the side and 2 inches below the planted seed. Starter band application of P (along with N) does not always lead to yield increases, but it can be an effective tool to enhance early season growth and reduce risks of losses associated with billbugs, competitive weeds, and summer droughts (see SoilFacts: Starter Phosphorus Fertilizer and Additives in North Carolina Soils: Use, Placement, and Plant Response, <http://www.soil.ncsu.edu/publications/Soilfacts/AG-439-75.pdf>). Use of starter fertilizers is more important in no-till since soil warming is delayed, and the cooler temperatures can reduce the rate of crop growth. For fields that already have very high soil test P levels (P-index >100), numerous North Carolina

tests have shown there is no advantage to applying additional P in a starter band. Even most mineral soils testing greater than a 50 P-Index generally do not need starter P.

The risk of salt injury is also related to the amount of salt applied. Generally, a maximum rate of 80 lb per acre of nitrogen alone, K₂O alone, or a combination of nitrogen plus K₂O is suggested for "2 x 2" band placement. If side placement is not possible or practical, thoroughly mix a broadcast application into the soil. If broadcast, a preferred method would be to apply before the land is plowed.

For tobacco, place the band 3 to 5 inches from the transplants. This distance reduces the chance of placing plants in fertilizer bands. If side placement equipment is not available, place the fertilizer deep in the row so it will be 3 to 5 inches below the roots of the transplants.

Seedlings of small grain or plants such as clovers, grasses, and alfalfa respond very early to phosphate; consequently, it is usually desirable to place phosphate close to the seed as is done with the conventional grain drill. Remember that when you use very high rates of nitrogen- and potassium-bearing fertilizers for small-seeded plants, there is some danger of fertilizer injury. A general guideline is not to exceed the rate of 80 lb/acre of N, K, or N + K in a 2-inch-by-2-inch banded starter placement. When a greater rate is necessary, make a split application; broadcast part and apply the remainder in the row at seeding.

Livestock Manure Production Rates and Nutrient Content

D.A. CROUSE, T. J. SMYTH, AND C. R. CROZIER, Soil Science; S. SHAH, Biological & Agricultural Engineering; and B. R. CLEVELAND, NCDA&CS

The use of livestock and poultry manure as a crop fertilizer supplement has come full circle. Before the advent of inexpensive inorganic fertilizers after World War II, farmers routinely used manure to complement a good fertility program. Today, because of rising costs of commercial fertilizers and increasing emphasis on sound manure management to protect water quality, renewed interest has been focused on maximizing the fertilizer returns of organic manures.

Waste production and characteristics are influenced by several factors such as the following. Waste from open housing systems and manure storage areas is diluted by rainfall. Manure drying reduces the nitrogen content because of associated ammonia losses. The longer manure remains in a housing or storage area before removal, the more chance there is for nitrogen loss. Liquid manure storage pits or basins retain the urine and manure fluids, which can contain as much as 50 percent of the total nitrogen. Lagoon treatment reduces total nitrogen by 50 to 85 percent and converts as much as 90 percent of the phosphorus to forms that settle into the sludge rather than remain in lagoon effluent. The following tables give average manure and wastewater characteristics for various handling methods, but because of the variability in nutrient values, wastes must be sampled and analyzed within 60 days of application. Waste samples can be analyzed for 11 essential plant nutrients at a nominal fee by the N.C. Department of Agriculture and Consumer Services, Waste Analysis Lab, 1040 Mail Service Center, Raleigh, NC 27699-1040; (919) 733-2655. When using a private carrier to deliver the samples or dropping them off in person, use the physical address of 4300 Reedy Creek Road, Raleigh, NC 27607-6465.

It is important that land application of manures becomes an integral part of the overall soil-fertility management strategy. Decomposition and mineralization of the manure in soil release significant amounts of nutrients essential for crop growth. More

nitrogen is conserved when manure is incorporated into the soil. If left on the surface, up to 25 percent of the ammonia nitrogen can be lost within 2 days, and 75 percent or more can be lost within 1 month after application. The total nutrients in Tables 4-14 through 4-20 can be multiplied by the appropriate availability coefficients from Table 4-21 to approximate plant nutrients available during the first year of application. The results on the NCDA&CS Waste Analysis Report already take into account first year nutrient availability (see the heading Nutrients Available for First Crop) and report them in pounds per ton or pounds per 1,000 gallons.

To calculate nutrient availability in the manure for plant uptake during the first year after application using the NCDA&CS Waste Analysis Report, multiply the results reported in parts per million (ppm) by the appropriate coefficient in Table 4-21. If the manure is solid, also multiply the ppm by the percent dry matter (%DM) given on the report. Apply manure as close to the period of maximum plant demand for nutrients as possible.

Base manure application rates on the available portion of the nutrients; do not apply more than the receiver crop needs. Excessive amounts not only waste valuable nutrients but may result in surface and/or groundwater pollution. Use soil testing (<http://ncagr.gov/agronomi/sthome.htm>) to predict nutrient and lime requirements and plant analysis; (<http://ncagr.gov/agronomi/uyrplant.htm>) to monitor the nutritional status and effectiveness of the nutrient management program. Additional animal manure information is available in Soil Facts AG-439-4, *Swine Manure as a Fertilizer Source*; AG-439-5, *Poultry Manure as a Fertilizer Source*; and AG-439-28, *Dairy Manure as a Fertilizer Source*. Fact sheets are available at county Cooperative Extension centers or electronically at (<http://www.soil.ncsu.edu/about/publications.php#WasteManagement>).

TABLE 4-14. LIVESTOCK FRESH MANURE CHARACTERISTICS

Manure Source	Average Animal Weight (lb)	Feces and Urine Production ¹		Density (lb/ft ³)	Total Solids TS (%w.b.)	Nitrogen		Phosphorus P ₂ O ₅ (lb/ton)	Potassium K ₂ O (lb/ton)
		(lb/day)	(tons/yr)			Total (lb/ton)	NH ₄ -N (lb/ton)		
BEEF	800	48.5	8.3	61.4	14.7	13.4	3.9	7.3	8.9
BROILER	2	0.16	0.024	63.7	25.6	26.3	6.7	16.3	11.7
DAIRY	1,400	122.3	22.3	62.5	13.9	10.4	1.9	5.1	8.2
DUCK	3	0.33	0.050	62.4	27.0	28.0	8.2	23.4	17.2
GOAT	140	5.8	1.1	62.5	32.5	21.8	6.1	12.1	23.6
HORSE	1,000	50.3	9.2	61.1	29.6	12.1	2.4	6.3	12.0
LAYER	4	0.26	0.047	61.6	24.9	26.6	6.6	21.3	11.6
RABBIT	10	0.31	0.056	28.0	51.6	24.5	0.4	25.2	11.1
SHEEP	60	2.4	0.4	62.3	28.1	20.8	5.8	9.4	19.0
SWINE	135	11.1	1.9	61.5	10.3	12.3	7.5	9.3	8.8
TURKEY	15	0.68	0.112	63.5	25.3	28.0	8.1	24.4	12.1
VEAL	200	12.4	2.0	62.2	6.5	7.9	3.7	4.0	10.6

¹ As voided.**TABLE 4-15. LIVESTOCK LOT SURFACE SCRAPED MANURE CHARACTERISTICS**

Manure Source	Average Animal Weight (lb)	Manure Accumulation		Density (lb/ft ²)	Total Solids TS (%w.b.)	Nitrogen		Phosphorus P ₂ O ₅ (lb/ton)	Potassium K ₂ O (lb/ton)
		lb/day	tons/yr			Total lb/ton	NH ₄ -N lb/ton		
BEEF paved feed lot ³ unpaved feedlot ³	800	33.5	5.7	59.7	28.6	14.3	4.9	9.4	12.9
		12.7	2.2	60.3	58.9	25.1	4.7	17.8	21.6
DAIRY ¹	1,400	88.0	16.1	56.4	21.8	10.3	2.5	7.1	8.6
HORSE ⁵	1,000	32.9	6.0	14.0	49.7	12.8	1.9	9.7	13.9
LAYER undercage scraped ² highrise stored ⁴	4	0.16	0.028	62.4	35.2	28.3	14.0	31.7	19.5
		0.11	0.020	51.2	52.8	38.6	11.8	51.1	26.3
SWINE ¹	135	7.8	1.3	62.4	18.5	13.0	5.6	13.3	9.1

¹ Manure collected within 1 week.² Manure collected within 2 days.³ Manure collected after each group of cattle.⁴ Annual manure accumulation on unpaved surface.⁵ Stable manure and bedding on unpaved surface.

TABLE 4-16. POULTRY HOUSE LITTER CHARACTERISTICS

Manure Source	Average Animal Weight (lb)	Manure and Litter Accumulation		Density (lb/ft ²)	Total Solids TS (% w.b.)	Nitrogen		Phosphorus P ₂ O ₅ (lb/ton)	Potassium K ₂ O (lb/ton)
		lb/day	tons/yr			Total lb/ton	NH ₄ -N lb/ton		
BROILER									
Broiler ¹	2.0	0.041	0.0063	31.7	78.6	72.1	12.0	69.3	46.6
Roaster ¹	4.0	0.063	0.010	29.0	76.2	69.5	16.0	69.9	46.7
Breeder ¹	6.0	0.13	0.023	50.0	68.6	37.5	8.0	58.3	35.2
TURKEY									
Poult ²	2.5	0.048	0.0070	22.9	79.5	40.1	9.6	43.3	26.9
Grower hen ¹	10.0	0.16	0.02	32.3	73.2	55.6	12.1	63.3	39.9
Grower tom ¹	15.0	0.24	0.041	32.3	73.2	55.6	12.1	63.3	39.9
DUCK¹	3.0	0.17	0.026	500	37.0	17.0	3.7	21.4	12.7

¹ Based on annual cleanout after full production.

² Based on cleanout after each group of birds.

TABLE 4-17. POULTRY STOCKPILED LITTER CHARACTERISTICS

Manure Source	Average Animal Weight (lb)	Manure and Litter Accumulation		Density (lb/ft)	Total Solids (TS) (% w.b.)	Nitrogen		Phosphorus P ₂ O ₅ lb/ton	Potassium K ₂ O lb/ton
		lb/day	tons/yr			Total lb/ton	NH ₄ -N lb/ton		
BROILER¹	2	0.038	0.0057	33.1	60.6	32.7	6.9	76.7	32.0
TURKEY¹	15	0.22	0.037	24.1	61.2	32.1	5.5	69.6	30.1
DUCK¹	3	0.09	0.014	50.0	49.1	22.3	4.3	41.2	21.7

¹ Annual house accumulation removed to uncovered stockpile to be spread within 6 months.

TABLE 4-18. LIVESTOCK LIQUID MANURE SLURRY CHARACTERISTICS

Manure Source	Average Animal Weight (lb)	Manure Slurry Accumulation ¹ (gal/animal/day)	Density (lb/gal)	Total Solids TS (% w.b.)	Nitrogen		Phosphorus P ₂ O ₅ (lb/1,000 gal)	Potassium K ₂ O (lb/1,000 gal)
					Total (lb/1,000 gal)	NH ₄ -N (lb/1,000 gal)		
BEEF	800	6.8	8.3	12.0	35.0	14.6	22.6	31.6
DAIRY	1,400	22.5	8.3	7.0	22.5	9.2	13.8	20.0
LAYER	4	0.046	7.8	11.2	57.6	36.8	52.0	33.1
SWINE	135	2.3	8.4	5.1	26.7	16.8	18.9	15.2
VEAL	200	1.4	8.5	2.6	30.4	25.3	17.0	42.1

¹ Based on 6 to 12 months' accumulation of manure, excess water usage, storage surface rainfall surplus; does not include fresh water for flushing or lot runoff.

TABLE 4-19. LIVESTOCK ANAEROBIC LAGOON LIQUID CHARACTERISTICS

Manure Source	Average Animal Weight (lb)	Lagoon Liquid Accumulation ¹ (acre-inch/animal/year)	Density (lb/gal)	Total Solids TS (% w.b.)	Nitrogen		Phosphorus P ₂ O ₅ (lb/1000 gal)	Potassium K ₂ O (lb/1000 gal)
					Total (lb/1000 gal)	NH ₄ -N (lb/1000 gal)		
BEEF	800	0.13	8.3	0.59	3.4	2.3	1.8	4.9
DAIRY	1,400	0.41	8.3	0.52	4.9	3.2	2.8	6.5
LAYER	4	0.00095	8.3	0.49	6.6	5.6	1.7	10.3
SWINE	135	0.035	8.3	0.32	4.7	3.8	1.9	4.9
VEAL	200	0.032	8.3	—	2.1	1.3	0.4	3.0

¹ Based on accumulation of manure, excess water usage, and average annual lagoon surface rainfall surplus; does not include fresh water for flushing or lot runoff.

TABLE 4-20. LIVESTOCK ANAEROBIC LAGOON SLUDGE CHARACTERISTICS

Manure Source	Average Animal Weight (lb)	Lagoon Sludge Accumulation ¹ (gal/hd/day)	Density (lb/gal)	Total Solids TS (% w.b.)	Nitrogen		Phosphorus P ₂ O ₅ (lb/1,000 gal)	Potassium K ₂ O (lb/1,000 gal)
					Total (lb/1,000 gal)	NH ₃ -N (lb/1,000 gal)		
BEEF	800	2.3	8.3	11.4	38.2	17.1	58.9	14.6
DAIRY	1,400	5.9	8.3	7.3	19.2	6.2	41.8	9.2
LAYER	4	0.013	8.3	16.5	20.8	6.5	77.2	9.8
SWINE	135	0.1	8.9	10.0	24.4	5.9	52.6	6.5

¹ No manure solids removed before waste enters the lagoon.

TABLE 4-21. LIVESTOCK MANURE NUTRIENT FIRST-YEAR AVAILABILITY COEFFICIENTS

Manure Source	Application Method							
	Injection ¹		Soil Incorporation ²		Broadcast ³		Irrigation ⁴	
	Plant Nutrient							
	N	Other	N	Other	N	Other	N	Other
PAVED SURFACE SCRAPED MANURE	—	—		0.8		0.7	—	—
Dairy			0.6		0.4			
Beef			0.6		0.4			
Swine			0.6		0.4			
Sheep			0.6		0.4			
Goat			0.5		0.5			
Rabbit			0.6		0.4			
Layer								
UNPAVED SURFACE MANURE ACCUMULATION	—	—		0.8		0.7	—	—
Beef, feedlot			0.6		0.5			
Horse, stable			0.5		0.5			
Layer, deep pit			0.6		0.4			
POULTRY HOUSE LITTER	—	—	0.6	0.8	0.5	0.7	—	—
Broiler								
Broiler breeder								
Turkey								
Duck								
POULTRY STOCKPILED LITTER	—	—	0.6	0.8	0.5	0.7	—	—
Broiler								
Turkey								
Duck								
LIQUID MANURE SLURRY		0.8		0.8	0.4	0.7		0.7
Dairy	0.7		0.6				0.4	
Beef	0.7		0.6				0.4	
Veal	0.9		0.7				0.3	
Swine	0.8		0.7				0.3	
Layer	0.8		0.7				0.3	
ANAEROBIC LAGOON LIQUID		0.8		0.8	0.5	0.7	0.5	0.7
Dairy	0.8		0.7					
Beef	0.7		0.7					
Veal	0.8		0.7					
Swine	0.9		0.8					
Layer	0.9		0.8					
ANAEROBIC LAGOON SLUDGE		0.8	0.6	0.8	0.4	0.7	0.4	0.7
Dairy	0.7							
Beef	0.7							
Swine	0.6							
Layer	0.6							

¹ Manure injected directly into soil and covered immediately.

² Surface-spread manure plowed or disked into soil within 2 days.

³ Surface-spread manure uncovered for 1 month or longer.

⁴ Sprinkler irrigated liquid uncovered for 1 month or longer.

Use of Municipal and Industrial Biosolids

J. G. WHITE, and D. L. LINDBO, Soil Science

Many municipalities and industries apply residuals (a.k.a., biosolids) and by-products from wastewater treatment facilities to agricultural land. Most of these biosolids contain sufficient plant nutrients (N, P, K) and liming value to make this practice beneficial to agriculture. Some biosolids may contain heavy metals and other substances that could pose a hazard either to crops being grown, the livestock or humans that consume them, and/or to the environment. However, guidelines developed by the USDA and the EPA regulate biosolids application rates so that these potential hazards are avoided or minimized.

Any municipality or industry that wishes to apply biosolids to its own land or to private farmland must obtain a permit from the North Carolina Department of Environment and Natural Resources (DENR). To obtain a permit, the municipality or industry must submit sufficient data to show that the proposed sites are suitable for biosolid application or that the biosolid qualifies as a "clean biosolid." Some of the factors considered in site selection are topography; soil type; distance from streams, wells, and property lines; flood hazard, depth to bedrock and the water table. If the site is suitable, then loading rates are calculated based on biosolid composition, nutrient requirement of the crop, previous history of biosolids application, and soil pH. If a property owner wishes to apply biosolids to his own land, he/she, too, must obtain a permit from DENR.

Many biosolids are treated with materials to raise sludge pH to reduce pathogen concentrations and vector attraction. These biosolids often possess liming capabilities that should be identified before use. If ignored and land-applied as a nutrient source, over-liming may occur and negatively affect plant growth.

Since the preparation of permit applications is the responsibility of the municipality, industry, and land owner, and since the process is somewhat involved, we will not attempt to outline it here. Information on the permitting procedure may be obtained from one of the regional offices of DENR or from Extension SoilFacts AG-439-6, *Permit Guidelines for Application of Municipal Sludge on Agricultural Lands*, which is available from your county Cooperative Extension Center or Department of Communication Services, Box 7603, N.C. State University, Raleigh, NC 27695-7603.

Since most biosolids produced in North Carolina are suitable for use on agricultural land, we encourage interested farmers to contact their county Cooperative Extension Service center about possible sources of biosolids in their area. Table 10-22 shows average characteristics of several biosolids. The user is strongly advised to obtain actual, current analysis of any product that is to be land-applied. For more information, see SoilFacts AG-439-19 *Using Municipal Solid Waste Compost and SoilFacts AG-439-19 Land Application of Municipal Sludge — Advantages and Concerns*.

TABLE 4-22. CHARACTERISTICS OF SEVERAL MUNICIPAL AND INDUSTRIAL BIOSOLIDS FROM OR AVAILABLE IN NORTH CAROLINA¹

Biosolid Source	Plant-available Nitrogen ²	P ₂ O ₅	K ₂ O	Ca	Mg	Cu	Zn	Ni	Cr	Pb	Cd	Mo	As
	Constituents in pounds per dry ton of sludge												
Milorganite	65	68	11.5	-	-	0.44	0.94	0.46	0.026	0.134	0.009	0.022	0.007
Mount Olive	43	46	12.5	8	6.6	0.48	1.05	0.029	0.067	0.085	0.005	0.016	0.004
Orange County	58	92	-	-	-	1.24	1.47	0.074	0.106	0.102	0.004	0.030	0.004
Elizabeth City	15	47	-	-	-	0.51	0.9	0.018	0.038	0.08	0.006	0.038	0.004
Charlotte Utilities													
Irwin Creek	16	49	-	-	-	0.63	0.9	0.098	0.230	0.210	0.006	0.10	0.062
Mallard Creek	35	69	-	-	-	0.28	0.84	0.036	0.044	0.042	0.004	0.022	0.003
McDowell Creek	36	133	-	-	-	0.27	0.78	0.032	0.078	0.054	0.003	0.028	0.005
Wilmington	31	103	2.6	-	-	1.0	19.0	0.05	0.10	0.082	0.007	0.022	0.010
Raleigh	32	114	46	-	-	0.58	1.7	0.082	0.024	0.056	0.007	0.028	0.001
Albemarle	9.5	3.8	16	-	-	0.38	0.42	0.140	0.136	0.005	0.006	0.026	0.024
High Point	24	96	4.3	-	-	0.40	0.62	0.034	0.082	0.094	0.004	0.032	0.005
Burlington Mills (industrial sludge)	35	52	-	-	-	0.122	0.54	0.044	0.94	0.021	0.002	0.01	0.006
Cone Mills (industrial sludge)	48	36	26	-	-	0.164	0.36	0.036	0.166	0.020	0.002	0.01	0.004

¹ Results between samples may vary. This table represents average or median values. Data from NC-DENR-Division of Water Quality.

² Plant-available nitrogen (PAN) is dependent on sludge processing methods and land application method.

